

## UBASH3A Polyclonal Antibody

**Catalog No.** E-AB-52430

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

|                     |                                                         |
|---------------------|---------------------------------------------------------|
| <b>Reactivity</b>   | Human, Mouse                                            |
| <b>Immunogen</b>    | Fusion protein of human UBASH3A                         |
| <b>Host</b>         | Rabbit                                                  |
| <b>Isotype</b>      | IgG                                                     |
| <b>Purification</b> | Antigen affinity purification                           |
| <b>Conjugation</b>  | Unconjugated                                            |
| <b>Buffer</b>       | PBS with 0.05% NaN <sub>3</sub> and 40% Glycerol, pH7.4 |


### Applications Recommended Dilution

**IHC** 1:40-1:250

### Data



Immunohistochemistry of paraffin-embedded Human tonsil tissue using UBASH3A Polyclonal Antibody at dilution of 1:50(×200)



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using UBASH3A Polyclonal Antibody at dilution of 1:50(×200)

### Preparation & Storage

**Storage** Store at -20°C. Avoid freeze / thaw cycles.

### Background

This gene encodes one of two family members belonging to the T-cell ubiquitin ligand (TULA) family. Both family members can negatively regulate T-cell signaling. This family member can facilitate growth factor withdrawal-induced apoptosis in T cells, which may occur via its interaction with AIF, an apoptosis-inducing factor. Alternative splicing of this gene results in multiple transcript variants.

### For Research Use Only