

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human, Mouse
Immunogen	Fusion protein of human UNC45B
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Formulation	PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

Applications Recommended Dilution


WB	1:500-1:2000
IHC	1:50-1:300
ELISA	1:5000-1:10000

Data



Western blot analysis of Human heart tissue lysate using UNC45B Polyclonal Antibody at dilution of 1:600

Observed Mw:Refer to figures
Calculated Mw:104/95 kDa



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using UNC45B Polyclonal Antibody at dilution of 1:65(×200)

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

This gene encodes a co-chaperone required for folding and accumulation of type II myosins. The protein consists of three tetratricopeptide repeat motifs at the N-terminus that form a complex with heat shock protein 90, a central region of unknown function that is conserved in all Unc-45 proteins, and a C-terminal Unc-45/Cro1/She4 domain. The protein is expressed at high levels in striated muscle, where its muscle myosin chaperone activity is dependent on heat shock protein 90 acting as a co-chaperone. A missense mutation in this gene has been associated with cataract development. Alternative splicing results in multiple transcript variants.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017