

NKD2 Polyclonal Antibody

Catalog No. E-AB-52880

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Immunogen	Fusion protein of human NKD2
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% NaN ₃ and 40% Glycerol,pH7.4

Applications

**WB 1:500-1:2000,
IHC 1:50-1:300,
ELISA
1:5000-1:10000**

Recommended Dilution


Data



Western blot analysis of A549 cell lysate using NKD2 Polyclonal Antibody at dilution of 1:600
Observed Mw:Refer to figures
Calculated Mw:50 kDa



Immunohistochemistry of paraffin-embedded Human tonsil tissue using NKD2 Polyclonal Antibody at dilution of 1:50(×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using NKD2 Polyclonal Antibody at dilution of 1:50(×200)

Preparation & Storage

For Research Use Only

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

This gene encodes a member of a family of proteins that function as negative regulators of Wnt receptor signaling through interaction with Dishevelled family members. The encoded protein participates in the delivery of transforming growth factor alpha-containing vesicles to the cell membrane. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.