

SLC16A7 Polyclonal Antibody

Catalog No. E-AB-53265

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Immunogen	Synthetic peptide of human SLC16A7
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% NaN ₃ and 40% Glycerol, pH7.4

Applications


Recommended Dilution

WB	1:500-1:2000
IHC	1:30-1:150

Data



Western blot analysis of Human abdominal interstitial sarcoma tissue lysate using SLC16A7 Polyclonal Antibody at dilution of 1:400
Observed Mw: Refer to figures
Calculated Mw: 52 kDa



Immunohistochemistry of paraffin-embedded Human tonsil tissue using SLC16A7 Polyclonal Antibody at dilution of 1:35 (×200)

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

This gene is a member of the monocarboxylate transporter family. Members in this family transport metabolites, such as lactate, pyruvate, and ketone bodies. The protein encoded by this gene catalyzes the proton-linked transport of monocarboxylates and has the highest affinity for pyruvate. This protein has been reported to be more highly expressed in prostate and colorectal cancer specimens when compared to control specimens. Alternative splicing results in multiple transcript variants.

For Research Use Only