

## CMKLR1 Polyclonal Antibody

Catalog No. E-AB-30909

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

|                     |  |
|---------------------|--|
| <b>Reactivity</b>   | Human,Mouse,Rat  |
| <b>Immunogen</b>    | Synthesized peptide derived from the Internal region of human ChemR23        |
| <b>Host</b>         | Rabbit   |
| <b>Isotype</b>      | IgG  |
| <b>Purification</b> | Affinity purification  |
| <b>Conjugation</b>  | Unconjugated   |
| <b>Buffer</b>       | PBS with 0.02% sodium azide, 0.5% protective protein and 50% glycerol, pH7.4 |

### Applications Recommended Dilution

|              |              |
|--------------|--------------|
| <b>WB</b>    | 1:500-1:2000 |
| <b>IHC</b>   | 1:100-1:300  |
| <b>IF</b>    | 1:200-1:1000 |
| <b>ELISA</b> | 1:20000      |

### Data



Western Blot analysis of 3T3, K562 cells using CMKLR1 Polyclonal Antibody at dilution of 1:1000.

**Observed Mw:42kDa**  
**Calculated Mw:42kDa**

### Preparation & Storage

**Storage** Store at -20°C. Avoid freeze / thaw cycles.

### Background

Receptor for the chemoattractant adipokine chemerin/RARRES2 and for the omega-3 fatty acid derived molecule resolvin E1. Interaction with RARRES2 induces activation of intracellular signaling molecules, such as SKY, MAPK1/3 (ERK1/2), MAPK14/P38MAPK and PI3K leading to multifunctional effects, like, reduction of immune responses, enhancing of adipogenesis and angiogenesis. Resolvin E1 down-regulates cytokine production in macrophages by reducing the activation of MAPK1/3 (ERK1/2) and NF-kappa-B. Positively regulates adipogenesis and adipocyte metabolism. Acts as a coreceptor for several SIV strains (SIVMAC316, SIVMAC239, SIVMACL7E-FR and SIVSM62A), as well as a primary HIV-1 strain (92UG024-2).

### For Research Use Only