

## CEACAM1 Polyclonal Antibody

**Catalog No.** E-AB-40352

*Note:* Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Reactivity</b>	Mouse,Rat
<b>Immunogen</b>	Recombinant Mouse Carcinoembryonic antigen-related cell adhesion molecule 1 protein
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Antigen Affinity Purification
<b>Conjugation</b>	Unconjugated
<b>Buffer</b>	PBS with 0.05% Proclin300, 50% glycerol, pH7.3.


### Applications Recommended Dilution

**IHC** 1:100-1:300


### Data



Immunohistochemistry of paraffin-embedded Mouse colon using Ceacam1 Polyclonal Antibody at dilution of 1:200



Immunohistochemistry of paraffin-embedded Mouse liver using Ceacam1 Polyclonal Antibody at dilution of 1:200



Immunohistochemistry of paraffin-embedded Rat liver using Ceacam1 Polyclonal Antibody at dilution of 1:200

### Preparation & Storage

**Storage** Store at -20°C. Avoid freeze / thaw cycles.

### Background

### For Research Use Only

This gene encodes a member of the carcinoembryonic antigen (CEA) gene family, which belongs to the immunoglobulin superfamily. Two subgroups of the CEA family, the CEA cell adhesion molecules and the pregnancy-specific glycoproteins, are located within a 1.2 Mb cluster on the long arm of chromosome 19. Eleven pseudogenes of the CEA cell adhesion molecule subgroup are also found in the cluster. The encoded protein was originally described in bile ducts of liver as biliary glycoprotein. Subsequently, it was found to be a cell-cell adhesion molecule detected on leukocytes, epithelia, and endothelia. The encoded protein mediates cell adhesion via homophilic as well as heterophilic binding to other proteins of the subgroup.