

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|---------------------|-----------------------------------------------------|
| Reactivity | Human,Mouse |
| Immunogen | Recombinant protein of human APPL1 |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Affinity purification |
| Conjugation | Unconjugated |
| Formulation | PBS with 0.05% sodium azide and 50% glycerol, PH7.4 |


Applications Recommended Dilution

| | |
|------------|--------------|
| WB | 1:500-1:2000 |
| IHC | 1:100-1:300 |


Data



Western Blot analysis of A172 cell using APPL1 Polyclonal Antibody at dilution of 1:600
Calculated Mw:80kDa



Immunohistochemistry of paraffin-embedded Human esophagus cancer using APPL1 Polyclonal Antibody at dilution of 1:70



Immunohistochemistry of paraffin-embedded Human lung cancer using APPL1 Polyclonal Antibody at dilution of 1:70

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

The protein encoded by this gene has been shown to be involved in the regulation of cell proliferation, and in the crosstalk between the adiponectin signalling and insulin signalling pathways. The encoded protein binds many other proteins, including RAB5A, DCC, AKT2, PIK3CA, adiponectin receptors, and proteins of the NuRD/MeCP1 complex. This protein is found associated with endosomal membranes, but can be released by EGF and translocated to the nucleus.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017