
MKP-3 Polyclonal Antibody

Catalog No. E-AB-10308

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|---|
| Reactivity | Human,Mouse,Rat |
| Immunogen | Recombinant protein of human DUSP6 |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Affinity purification |
| Conjugation | Unconjugated |
| Buffer | PBS with 0.05% sodium azide and 50% glycerol, PH7.4 |
| Applications | Recommended Dilution |
| WB | 1:500-1:2000 |
| Data | |



Western Blot analysis of Jurkat cell and Human fetal kidney tissue using MKP-3 Polyclonal Antibody at dilution of 1:450 Calculated Mw:27kDa

Preparation & Storage

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Background

The protein encoded by this gene is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product inactivates ERK2, is expressed in a variety of tissues with the highest levels in heart and pancreas, and unlike most other members of this family, is localized in the cytoplasm. Two transcript variants encoding different isoforms have been found for this gene.