
CAPG Polyclonal Antibody

Catalog Number:E-AB-10720



Note: Centrifuge before opening to ensure complete recovery of vial contents.


| Description | |
|--------------|---|
| Reactivity | Human,Mouse,Rat |
| Immunogen | Recombinant protein of human CAPG |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Affinity purification |
| Conjugation | Unconjugated |
| Formulation | PBS with 0.05% sodium azide and 50% glycerol, PH7.4 |
| Applications | Recommended Dilution |
| WB | 1:1000-1:5000 |
| IHC | 1:50-1:200 |
| Data | |





Immunohistochemistry of paraffin-embedded Human brain using CAPG Polyclonal Antibody at dilution of 1:40

Western Blot analysis of Jurkat and RAW264.7 cell using CAPG Polyclonal Antibody at dilution of 1:950 Calculated Mw:38kDa



Immunohistochemistry of paraffin-embedded Human esophagus cancer using CAPG Polyclonal Antibody at dilution of 1:40

Preparation & Storage

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Background

This gene encodes a member of the gelsolin/villin family of actin-regulatory proteins. The encoded protein reversibly blocks the barbed ends of F-actin filaments in a Ca2+ and phosphoinositide-regulated manner, but does not sever preformed actin filaments. By capping the barbed ends of actin filaments, the encoded protein contributes to the control of actin-based motility in non-muscle cells. Alternatively spliced transcript variants have been observed for this gene.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Web: www.elabscience.com Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>