## **KLC1** Polyclonal Antibody

Catalog No. E-AB-10932

*Note:* Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human,Mouse,Rat
Immunogen	Recombinant protein of human KLC1
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% sodium azide and 50% glycerol, PH7.4
Applications	Recommended Dilution
WB	1:200-1:1000
IHC	1:50-1:200
Data	



Western Blot analysis of Lovo cell using KLC1 Polyclonal Antibody at dilution of 1:500 Calculated Mw:65kDa



Immunohistochemistry of paraffin-embedded Human cervical cancer using KLC1 Polyclonal Antibody at dilution of 1:40



Immunohistochemistry of paraffin-embedded Human brain using KLC1 Polyclonal Antibody at dilution of 1:40

## **Preparation & Storage**

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Background

## **For Research Use Only**

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

## **Elabscience**®

Conventional kinesin is a tetrameric molecule composed of two heavy chains and two light chains, and transports various cargos along microtubules toward their plus ends. The heavy chains provide the motor activity, while the light chains bind to various cargos. This gene encodes a member of the kinesin light chain family. It associates with kinesin heavy chain through an N-terminal domain, and six tetratricopeptide repeat (TPR) motifs are thought to be involved in binding of cargos such as vesicles, mitochondria, and the Golgi complex. Thus, kinesin light chains function as adapter molecules and not motors per se. Although previously named "kinesin 2", this gene is not a member of the kinesin-2 / kinesin heavy chain subfamily of kinesin motor proteins. Extensive alternative splicing produces isoforms with different C-termini that are proposed to bind to different cargos; however, the full-length nature and/or biological validity of most of these variants have not been determined.

For Research Use Only