# **ERCC6L Polyclonal Antibody**

Catalog Number: E-AB-11196



Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

**Reactivity** Human

Immunogen Recombinant protein of human ERCC6L

Host Rabbit
Isotype IgG


Purification Affinity purification
Conjugation Unconjugated

**Formulation** PBS with 0.05% sodium azide and 50% glycerol, PH7.4


**Applications** Recommended Dilution

IHC 1:50-1:200

## Data



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using ERCC6L Polyclonal Antibody at dilution 1:30



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using ERCC6L Polyclonal Antibody at dilution 1:30

### Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

#### **Background**

PICH (Plk1-interacting checkpoint helicase), also known as DNA excision repair protein ERCC-6-like (ERCC6L) or tumor antigen BJ-HCC-15, is a 1,250 amino acid protein belonging to the SNF2/RAD54 helicase family. PICH is a DNA helicase and an essential component of the spindle assembly checkpoint. During mitosis, PICH recruits MAD2 to kinetochores and also regulates the tension on centromic chromatin. PICH is concentrated in between the kinetochores in prometophase cells, while in metaphase it localizes to the thin threads composed of catenated centromeric DNA that stretch between sister kinetochores. PICH is phosphorylated by Plk, which prevents PICH from associating with chromosome arms and restricts the localization of PICH to the kinetochore-centromere region. PICH/Plk interaction is also required for correct Plk localization to the kinetochore. PICH contains one helicase ATP-binding domain, two TPR repeats and one helicase C-terminal domain.

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com