# CD42b Polyclonal Antibody

Catalog Number: E-AB-11281



Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Reactivity Human, Mouse

Immunogen Recombinant protein of human GP1BA

Host Rabbit
Isotype IgG


Purification Affinity purification
Conjugation Unconjugated

**Formulation** PBS with 0.05% sodium azide and 50% glycerol, PH7.4

## **Applications** Recommended Dilution


WB 1:500-1:2000 IHC 1:100-1:300

#### Data




Western Blot analysis of Mouse liver tissue using CD42b Polyclonal Antibody at dilution of 1:800

Calculated Mw:72kDa



Immunohistochemistry of paraffin-embedded Human liver cancer using CD42b Polyclonal Antibody at dilution of 1:60



Immunohistochemistry of paraffin-embedded Spleen tissue using CD42b Polyclonal Antibody at dilution of 1:80(Elabscience® Product Detected by Lifespan).

## **Preparation & Storage**

Storage Store at -20°C. Avoid freeze / thaw cycles.

### Background

Glycoprotein Ib (GP Ib) is a platelet surface membrane glycoprotein composed of a heterodimer, an alpha chain and a beta chain, that is linked by disulfide bonds. The Gp Ib functions as a receptor for von Willebrand factor (VWF). The complete receptor complex includes noncovalent association of the alpha and beta subunits with platelet glycoprotein IX

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a>
Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>

# CD42b Polyclonal Antibody

Catalog Number: E-AB-11281



and platelet glycoprotein V. The binding of the GP Ib-IX-V complex to VWF facilitates initial platelet adhesion to vascular subendothelium after vascular injury, and also initiates signaling events within the platelet that lead to enhanced platelet activation, thrombosis, and hemostasis. This gene encodes the alpha subunit. Mutations in this gene result in Bernard-Soulier syndromes and platelet-type von Willebrand disease. The coding region of this gene is known to contain a polymophic variable number tandem repeat (VNTR) domain that is associated with susceptibility to nonarteritic anterior ischemic optic neuropathy.

### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>