
IFT74 Polyclonal Antibody

Catalog Number: E-AB-13159



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|---|
| Reactivity | Human |
| Immunogen | Synthetic peptide of human IFT74 |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Affinity purification |
| Conjugation | Unconjugated |
| Formulation | PBS with 0.05% sodium azide and 50% glycerol, PH7.4 |
| Applications | Recommended Dilution |
| WB | 1:500-1:2000 |
| Data | |



Western Blot analysis of HepG2 and 293T cell using IFT74 Polyclonal Antibody at dilution of 1:1400 Calculated Mw:69kDa

Preparation & Storage

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Background

Capillary morphogenesis protein 1 (CMG1), also known as intraflagellar transport protein 74 (ITP74) or coiled-coil domain-containing protein 2 (CCDC2), is a 600 amino acid human homologue of IFT-71, a complex B protein supporting intraflagellar transport (IFT) in Chlamydomonas. CMG1 localizes to the cytoplasmic vesicle and is highly expressed in adult and fetal kidney and testis, with lower levels of expression in adult heart, placenta, lung, liver and pancreas, and in fetal heart, lung and liver. CMG1 has been suggested to have a role in the primary cilia of HUVEC, and it also functions as a transcriptional regulator of cyclin D2 in spermatocyte-derived cells.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com