

RAGE Polyclonal Antibody

Catalog No. E-AB-13783

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human
Immunogen	Recombinant protein of human AGER
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% sodium azide and 50% glycerol, PH7.4


Applications Recommended Dilution

WB	1:500-1:2000
IHC	1:100-1:300


Data



Western Blot analysis of 231 cell using RAGE Polyclonal Antibody at dilution of 1:750
Calculated Mw:43kDa



Immunohistochemistry of paraffin-embedded Human breast cancer using RAGE Polyclonal Antibody at dilution of 1:70



Immunohistochemistry of paraffin-embedded Human gastric cancer using RAGE Polyclonal Antibody at dilution of 1:70

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

For Research Use Only

The advanced glycosylation end product (AGE) receptor encoded by this gene is a member of the immunoglobulin superfamily of cell surface receptors. It is a multiligand receptor, and besides AGE, interacts with other molecules implicated in homeostasis, development, and inflammation, and certain diseases, such as diabetes and Alzheimer's disease. Many alternatively spliced transcript variants encoding different isoforms, as well as non-protein-coding variants, have been described for this gene.