

TAC1 Polyclonal Antibody

Catalog No. E-AB-14247

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human,Mouse,Rat
Immunogen	Recombinant protein of human TAC1
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% sodium azide and 50% glycerol, PH7.4


Applications Recommended Dilution

IHC	1:25-1:100
------------	------------


Data



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TAC1 Polyclonal Antibody at dilution 1:25



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TAC1 Polyclonal Antibody at dilution 1:25



Immunohistochemistry of paraffin-embedded Colon, Submucosal Plexus tissue using TAC1 Polyclonal Antibody at dilution of 1:100(Elabscience® Product Detected by Lifespan).

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

For Research Use Only

This gene encodes four products of the tachykinin peptide hormone family, substance P and neurokinin A, as well as the related peptides, neuropeptide K and neuropeptide gamma. These hormones are thought to function as neurotransmitters which interact with nerve receptors and smooth muscle cells. They are known to induce behavioral responses and function as vasodilators and secretagogues. Multiple transcript variants encoding different isoforms have been found for this gene.