

CCDC99 Polyclonal Antibody

Catalog No. E-AB-14370

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human,Mouse
Immunogen	Recombinant protein of human SPDL1
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% sodium azide and 50% glycerol, PH7.4

Applications

Recommended Dilution

WB 1:200-1:1000

Data



Western Blot analysis of Mouse brain tissue using
CCDC99 Polyclonal Antibody at dilution of 1:300
Calculated Mw:70kDa

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Required for the localization of dynein and dynactin to the mitotic kintochore. Dynein is believed to control the initial lateral interaction between the kinetochore and spindle microtubules and to facilitate the subsequent formation of end-on kinetochore-microtubule attachments mediated by the NDC80 complex. Also required for correct spindle orientation. Does not appear to be required for the removal of spindle assembly checkpoint (SAC) proteins from the kinetochore upon bipolar spindle attachment. Interacts with KNTC1 and ZW10. These interactions appear weak and may be transient or indirect.

For Research Use Only