

ACOX3 Polyclonal Antibody

Catalog No. E-AB-14480

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|---------------------|---|
| Reactivity | Human |
| Immunogen | Recombinant protein of human ACOX3 |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Affinity purification |
| Conjugation | Unconjugated |
| Buffer | PBS with 0.05% sodium azide and 50% glycerol, PH7.4 |


Applications Recommended Dilution

| | |
|------------|------------|
| IHC | 1:50-1:200 |
|------------|------------|

Data



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using ACOX3 Polyclonal Antibody at dilution 1:35



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using ACOX3 Polyclonal Antibody at dilution 1:35

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Acyl-Coenzyme A oxidase 3 also known as pristanoyl -CoA oxidase (ACOX3) is involved in the desaturation of 2-methyl branched fatty acids in peroxisomes. Unlike the rat homolog, the human gene is expressed in very low amounts in liver such that its mRNA was undetectable by routine Northern-blot analysis or its product by immunoblotting or by enzyme activity measurements. However the human cDNA encoding a 700 amino acid protein with a peroxisomal targeting C-terminal tripeptide S-K-L was isolated and is thought to be expressed under special conditions such as specific developmental stages or in a tissue specific manner in tissues that have not yet been examined.

For Research Use Only