

HSP40 Polyclonal Antibody

Catalog No. E-AB-14642

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human, Mouse
Immunogen	Recombinant protein of human DNAJB1
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% sodium azide and 50% glycerol, PH7.4


Applications Recommended Dilution

WB	1:500-1:2000
IHC	1:50-1:200


Data



Western Blot analysis of 293T cell using HSP40 Polyclonal Antibody at dilution of 1:750
Calculated Mw:38kDa



Immunohistochemistry of paraffin-embedded Human lung cancer using HSP40 Polyclonal Antibody at dilution of 1:40



Immunohistochemistry of paraffin-embedded Human gastric cancer using HSP40 Polyclonal Antibody at dilution of 1:40

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

For Research Use Only

Heat shock protein 40 (HSP 40) family proteins bind to HSP 70 through their J-domain and regulate the function of HSP 70 by stimulating HSP 70 ATPase activity. HSP 40, also known as DnaJ, functions together with DnaK (HSP 70) and GrpE as a molecular chaperone, involving them in assembly and disassembly of protein complexes, protein folding, renaturation of denatured proteins, prevention of protein aggregation and protein export. HSP 40 stimulates the association between HSC 70 and HIP and translocates rapidly from the cytoplasm to the nuclei, and especially to the nucleoli, upon heat shock. There are five known HSP 40 family proteins.