

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human,Rat
Immunogen	Synthetic peptide of human TXN
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Formulation	PBS with 0.05% sodium azide and 50% glycerol, PH7.4


Applications Recommended Dilution

WB	1:1000-1:5000
IHC	1:50-1:200


Data



Western Blot analysis of Hela and MCF7 cell,
Human liver cancer tissue using TXN Polyclonal
Antibody at dilution of 1:600
Calculated Mw:12kDa



Immunohistochemistry of paraffin-embedded
Human thyroid cancer using TXN Polyclonal
Antibody at dilution of 1:40



Immunohistochemistry of paraffin-embedded
Human lung cancer using TXN Polyclonal Antibody
at dilution of 1:40

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

The protein encoded by this gene acts as a homodimer and is involved in many redox reactions. The encoded protein is active in the reversible S-nitrosylation of cysteines in certain proteins, which is part of the response to intracellular nitric oxide. This protein is found in the cytoplasm. Two transcript variants encoding different isoforms have been found for

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

TXN Polyclonal Antibody

Catalog Number:E-AB-16096



this gene. Plays a role in the reversible S-nitrosylation of cysteine residues in target proteins, and thereby contributes to the response to intracellular nitric oxide. Nitrosylates the active site Cys of CASP3 in response to nitric oxide (NO), and thereby inhibits caspase-3 activity.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017