

SLC7A11 Polyclonal Antibody

Catalog No. E-AB-17625

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human, Mouse
Immunogen	Synthetic peptide of human SLC7A11
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% sodium azide and 50% glycerol, PH7.4

Applications Recommended Dilution


IHC	1:20-100
------------	----------

Data



Immunohistochemistry of paraffin-embedded Human brain using SLC7A11 Polyclonal Antibody at dilution of 1/35

Calculated Mw:55 kDa



Immunohistochemistry of paraffin-embedded Human thyroid cancer using SLC7A11 Polyclonal Antibody at dilution of 1/35

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

This gene encodes a member of a heteromeric, sodium-independent, anionic amino acid transport system that is highly specific for cysteine and glutamate. In this system, designated Xc(-), the anionic form of cysteine is transported in exchange for glutamate. This protein has been identified as the predominant mediator of Kaposi sarcoma-associated herpesvirus fusion and entry permissiveness into cells. Also, increased expression of this gene in primary gliomas (compared to normal brain tissue) was associated with increased glutamate secretion via the XCT channels, resulting in neuronal cell death.

For Research Use Only