Recombinant Human IL-8/CXCL8 Protein (HEK293 Cells)

Catalog Number: PKSH030275



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Synonyms	Interleukin-8;IL-8;C-X-C Motif Chemokine 8;Emoctakin;Granulocyte Chemotactic Protein 1;GCP-1;Monocyte-Derived Neutrophil Chemotactic Factor;MDNCF;Monocyte-Derived Neutrophil-Activating Peptide;MONAP;Neutrophil-Activating Protein 1;NAP-1;Protein 3-10C;T-Cell Chemotactic Factor;GCP1;IL8;Interleukin-8;LECT;LUCT;LYNAP;MDNCF;MON AP;NAF;NAP-1;NAP1		
Species	Human		
Expression Host	HEK293 Cells		
Sequence	Ser 28-Ser 99		
Accession	NP_000575.1		
Calculated Molecular Weight	8.5 kDa		
Observed molecular weight	12 kDa		
Tag	None		
Properties			
Purity	> 95 % as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.		
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from sterile PBS, pH 7.5, 5% trehalose, 5% mannitol, 0.01% tween-80 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printe		
Reconstitution	Please refer to the printed manual for detailed information.		
Data			

	KDa	MK	R
	116	Ctable C	
Elabscie	66.2	-	olence
Elabo	45.0	-	Elabsoit
	35.0	-	
Elabscie	25.0	-	Elabscier
	18.4	dence	
	14.4	-	

> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin 8 (IL-8), also known as CXCL8, which is a chemokine with a defining CXC amino acid motif that was initially characterized for its leukocyte chemotactic activity, is now known to possess tumorigenic and proangiogenic properties as well. This chemokine is secreted by a variety of cell types including monocyte/macrophages, T cells,

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Catalog Number: PKSH030275



neutrophils, fibroblasts, endothelial cells, and various tumor cell lines in response to inflammatory stimuli. In human gliomas, IL-8 is expressed and secreted at high levels both in vitro and in vivo, and recent experiments suggest it is critical to glial tumor neovascularity and progression. Levels of IL-8 correlate with histologic grade in glial neoplasms, and the most malignant form, glioblastoma, shows the highest expression in pseudopalisading cells around necrosis, suggesting that hypoxia/anoxia may stimulate expression. Accumulating evidence has demonstrated that various types of cells can produce a large amount of IL-8/CXCL8 in response to a wide variety of stimuli, including proinflammatory cytokines, microbes and their products, and environmental chang. Numerous observations have established IL-8/CXCL8 as a key mediator in neutrophil-mediated acute inflammation due to its potent actions on neutrophils. The discovery of these biological functions suggests that IL-8/CXCL8 has crucial roles in various pathological conditions such as chronic inflammation and cancer.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine
Toll-free: 1-888-852-8623 Tel: 1-832-243-6086
Web: www.elabscience.com Email: techsupport@elabscience.com