

## Recombinant Human STK16/PKL12/MPSK Protein (His & NusA Tag)

Catalog No. PKSH030340

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	KRCT;MPSK;PKL12;TSF1
<b>Species</b>	Human
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Met 1-Ile 305
<b>Accession</b>	AAH02618.1
<b>Calculated Molecular Weight</b>	92.0 kDa
<b>Observed molecular weight</b>	105 kDa
<b>Tag</b>	N-His-NusA
<b>Bioactivity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 85 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
<b>Shipping</b>	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
<b>Formulation</b>	Supplied as sterile solution of PBS, pH 7.4
<b>Reconstitution</b>	Not Applicable

### Data



> 85 % as determined by reducing SDS-PAGE.

### Background

Serine/threonine-protein kinase 16, also known as myristoylated and palmitoylated serine/threonine-protein kinase, Protein kinase PKL12, TGF-beta-stimulated factor 1, TSF-1, MPSK1 and STK16, is a membrane protein which is ubiquitously expressed at very low levels. STK16 / MPSK1 belongs to the protein kinase superfamily and Ser/Thr protein kinase family. It contains one protein kinase domain. Transforming growth factor-beta (TGF-beta) shows a variety of biological activities in various organs or cells. Some factors such as Smads and TGF-beta activating kinase 1 have been characterized as signalling molecules downstream of TGF-beta. Several TGF-beta response elements have been identified

### For Research Use Only

such as cAMP response element, Smad binding element, and recognition sites for activating protein-1 and stimulating protein-1 in various gene promoters. STK16 / MPSK1 is an unique factor with two biological functions, transcriptional regulation and protein phosphorylation, that may be involved in TGF-beta signals. STK16 / MPSK1 is a protein kinase that act on both serine and threonine residues. STK16 / MPSK1 possessed DNA-binding ability and activated the TGF-beta responsive CNP promoter or vascular endothelial growth factor gene promoter which possesses a sequence element analogous to the TGF-beta responsive GC-rich element of the CNP promoter. STK16 / MPSK1 did not directly activate a Smads-dependent promoter from plasminogen activator inhibitor 1 gene, but it showed enhancement in co-operation with Smad3 and Smad4. STK16 / MPSK1 mRNA as well as its protein level were stimulated by TGF-beta treatment.