A Reliable Research Partner in Life Science and Medicine

Recombinant Human MAP4K2/GC Kinase Protein (His & GST Tag)

Catalog No. PKSH030347

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms BL44;GCK;MAP4K2;RAB8IP

Species Human

Expression Host Baculovirus-Insect Cells

SequenceMet 1-Tyr 812AccessionAAH47865.1Calculated Molecular Weight119 kDaObserved molecular weight116 kDaTagN-His-GST

Bioactivity Not validated for activity

Properties

Purity > 87 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.


Shipping This product is provided as liquid. It is shipped at frozen temperature with blue

ice/gel packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as sterile solution of 50mM Tris, 100mM NaCl, pH 8.0

Reconstitution Not Applicable

Data



> 87 % as determined by reducing SDS-PAGE.

Background

Mitogen-activated protein kinase kinase kinase 2, also known as B lymphocyte serine/threonine-protein kinase, Germinal center kinase, MAPK/ERK kinase kinase 2, MEK kinase kinase 2, Rab8-interacting protein and MAP4K2, is a cytoplasm and peripheral membrane protein which belongs to the protein kinase superfamily, STE Ser/Thr protein kinase family and STE20 subfamily. MAP4K2 contains one CNH domain and one protein kinase domain. Although this kinase is found in many tissues, its expression in lymphoid follicles is restricted to the cells of germinal centre, where it may participate in B-cell differentiation. MAP4K2 can be activated by TNF-alpha, and has been shown to

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com





A Reliable Research Partner in Life Science and Medicine

specifically activate MAP kinases. It is also found to interact with TNF receptor-associated factor 2 (TRAF2), which is involved in the activation of MAP3K1 / MEKK1. MAP4K2 enhances MAP3K1 oligomerization, which may relieve amino-terminal mediated MAP3K1 autoinhibition and lead to activation following autophosphorylation. It may also play a role in the regulation of vesicle targeting or fusion.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com