

## Recombinant Human DDR1 Kinase/MCK10 Protein (aa 444-913, His & GST Tag)

Catalog No. PKSH030389

*Note:* Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	CAK;CD167;DDR;EDDR1;HGK2;MCK10;NEP;NTRK4;PTK3;PTK3A;RTK6;TRKE
<b>Species</b>	Human
<b>Expression Host</b>	Baculovirus-Insect Cells
<b>Sequence</b>	Arg444-Val913
<b>Accession</b>	Q08345-1
<b>Calculated Molecular Weight</b>	80.0 kDa
<b>Observed molecular weight</b>	80 kDa
<b>Tag</b>	N-His-GST
<b>Bioactivity</b>	The specific activity was determined to be 2.75 nmol/min/mg using synthetic AXLtide peptide(CKKSRGDYMTMQIG) as substrate.

### Properties

<b>Purity</b>	> 89 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
<b>Shipping</b>	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
<b>Formulation</b>	Supplied as sterile solution of 20mM Tris, 500mM NaCl, pH 7.4, 10% glycerol, 3mM DTT
<b>Reconstitution</b>	Not Applicable

### Data



> 89 % as determined by reducing SDS-PAGE.

### Background

Discoidin domain receptor family, member 1 (DDR1), also known as or CD167a (cluster of differentiation 167a), and

### For Research Use Only

Mammary carcinoma kinase 10 (MCK10), belongs to a subfamily of tyrosine kinase receptors with an extracellular domain homologous to Dictyostellium discoideum protein discoidin 1. Receptor tyrosine kinases play a key role in the communication of cells with their microenvironment. These kinases are involved in the regulation of cell growth, differentiation and metabolism. Expression of DDR1/MCK10/CD167 is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, it has been shown to be significantly overexpressed in several human tumors. DDR1/MCK10/CD167 plays an important role in regulating attachment to collagen, chemotaxis, proliferation, and MMP production in smooth muscle cells. DDR1 functions in a feedforward loop to increase p53 levels and at least some of its effectors. Inhibition of DDR1 function resulted in strikingly increased apoptosis of wild-type p53-containing cells in response to genotoxic stress through a caspase-dependent pathway.

---

## For Research Use Only

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017