

Recombinant Human MAPKAPK3 Protein (GST Tag)

Catalog No. PKSH030397

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	3PK;MAPKAP-K3;MAPKAP3;MAPKAPK-3;MK-3
Species	Human
Expression Host	Baculovirus-Insect Cells
Sequence	Met 1-Gln 382
Accession	NP_004626.1
Calculated Molecular Weight	69.0 kDa
Observed molecular weight	69 kDa
Tag	N-GST
Bioactivity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20°C.
Formulation	Supplied as sterile solution of 50mM Tris, 100mM NaCl, pH 7.5, 0.25mM DTT, 0.1mM EDTA, 0.5mM PMSF, 10% glycerol
Reconstitution	Not Applicable

Data



> 90 % as determined by reducing SDS-PAGE.

Background

The MAPKAP kinases are a group of MAP kinase substrates which are themselves kinases. In response to activation, the MAP kinases phosphorylate downstream components on a consensus Pro-X-Ser/Thr-Pro motif. Several kinases that contain this motif have been identified and serve as substrates for the ERK and p38 MAP kinases. Mitogen-activated protein (MAP) kinase-activated protein kinase 3, also known as MAPKAPK-3 and 3pK, is a member of the Ser/Thr protein kinase family. It is Widely expressed in human tissues, with a higher expression level observed in heart and

For Research Use Only

skeletal muscle. No expression in brain. MAPKAPK-3 is unique since it was shown to be activated by three members of the MAPK family, namely extracellular-signal-regulated kinase (ERK), p38, and Jun-N-terminal kinase (JNK). It is highly activated both by mitogens and by stress-inducing agents or proinflammatory cytokines, and translocates to the cytoplasm from nucleus. MAPKAPK-3 is exclusively activated via the classical MAPK cascade, while stress-induced activation of MAPKAPK-3 is mainly mediated by p38, however the mechanism defining the specificity remains unknown.

For Research Use Only