

Recombinant Human DDR2 Kinase/CD167b Protein (aa 422-855, His & GST Tag)

Catalog No. PKSH030420

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CD167;MIG20a;NTRKR3;TKT;TYRO10
Species	Human
Expression Host	Baculovirus-Insect Cells
Sequence	Arg422-Glu855
Accession	Q16832
Calculated Molecular Weight	77.1 kDa
Observed molecular weight	77 kDa
Tag	N-His-GST
Bioactivity	The specific activity was determined to be 8 nmol/min/mg using synthetic AXLtide peptide(CKKSRGDYMTMQIG) as substrate.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
Formulation	Supplied as sterile solution of 20mM Tris, 500mM NaCl, pH 7.4, 10% glycerol
Reconstitution	Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Discoidin domain receptor 2 (DDR2) or CD167b (cluster of differentiation 167b) is a kind of protein tyrosine kinases associated with cell proliferation and tumor metastasis, and collagen, identified as a ligand for DDR2, up-regulates matrix metalloproteinase 1 (MMP-1) and MMP-2 expression in cellular matrix. DDR2/CD167b was found to recognise the

For Research Use Only

triple-helical region of collagen X as well as the NC1 domain. Binding to the collagenous region was dependent on the triple-helical conformation. DDR2/CD167b autophosphorylation was induced by the collagen X triple-helical region but not the NC1 domain, indicating that the triple-helical region of collagen X contains a specific DDR2 binding site that is capable of receptor activation. DDR2/CD167b is induced during stellate cell activation and implicate the phosphorylated receptor as a mediator of MMP-2 release and growth stimulation in response to type I collagen. Moreover, type I collagen-dependent upregulation of DDR2/CD167b expression establishes a positive feedback loop in activated stellate cells, leading to further proliferation and enhanced invasive activity.