

Recombinant Human CD31/PECAM1 Protein

Catalog No. PKSH030474

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Platelet endothelial cell adhesion molecule;PECAM-1;EndoCAM;GPIIA;PECA1;CD31;PECAM1
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Lys 601
Accession	EAW94208.1
Calculated Molecular Weight	65.1 kDa
Observed molecular weight	115 kDa
Tag	None
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 100mM Glycine, 10mM NaCl, 50mM Tris, pH 7.5 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

The Cluster of Differentiation 31 (CD31) adhesion molecule, also known as platelet-endothelial cell adhesion molecule-1

For Research Use Only

(PECAM-1), is the only known member of the CAM family on platelets. CD31 protein is a 130-kDa transmembrane glycoprotein expressed by endothelial cells, platelets, monocytes, neutrophils, and certain T cell subsets. CD31 protein is also expressed in certain tumors, including epithelioid hemangioendothelioma, other vascular tumors, and histiocytic malignancies. CD31 plays a key role in removing aged neutrophils and tissue regeneration. CD31 protein mediates the homotypic or heterotypic cell adhesion by binding to itself or the leukocyte integrin $\alpha v \beta 3$, and thus plays a role in neutrophil recruitment in inflammatory responses, transendothelial migration of leukocytes, as well as in cardiovascular development.

For Research Use Only