# Recombinant Human FGFR3/CD333 Protein (alpha(IIIb), Fc Tag)



Catalog Number: PKSH030490

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Fibroblast growth factor receptor 3;FGFR-3;FGFR3;JTK4;IIIc;ACH;CD333;CEK2;HSFGFR3EX
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Gly377
Accession	NP_001156685.1
Calculated Molecular Weight	65.3 kDa
Tag	C-hFc
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per $\mu$ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	



>95 % as determined by reducing SDS-PAGE.

### Background

FGFR3, also known as CD333, is a member of the fibroblast growth factor receptor (FGFR) family, with its amino acid sequence being highly conserved between members and among divergent species. FGFR family members differ from one another in their ligand affinities and tissue distribution. FGFRs are transmembrane catalytic receptors that have intracellular tyrosine kinase activity. Mutations in FGFR genes are the cause of several human developmental disorders characterized by skeletal abnormalities such as achondroplasia, and upregulation of FGFR expression may lead to cell transformation and cancer. FGFR3, a full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of FGFR3 interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. FGFR3 binds acidic and basic fibroblast

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

## Recombinant Human FGFR3/CD333 Protein (alpha(IIIb), Fc Tag)



Catalog Number:PKSH030490

growth hormone and plays a role in bone development and maintenance. Mutations in FGFR3 gene lead to craniosynostosis and multiple types of skeletal dysplasia. Three alternatively spliced transcript variants that encode different protein isoforms have been described. CD333 is the receptor for acidic and basic fibroblast growth factors.

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com