

Recombinant Human TSPAN8/Tetraspanin 8 Protein (His Tag)

Catalog No. PKSH030523

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CO-029;TM4SF3
Species	Human
Expression Host	HEK293 Cells
Sequence	Lys110-Asn205
Accession	P19075
Calculated Molecular Weight	13.4 kDa
Observed molecular weight	17&13 kDa
Tag	N-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Tetraspanin 8 (TSPAN8) as an important modulator of melanoma invasiveness; and several of its transcriptional regulators; which affect TSPAN8 expression during melanoma progression toward an invasive stage. p53 as a negative

For Research Use Only

regulator of Tspan8 expression. p53 as a regulator of melanoma invasion and the concept that reactivating p53 could provide a strategy for modulating not only proliferative but also invasive capacity in melanoma treatment. Tetraspanin 8 (TSPAN8) is a tumor-associated antigen implicated in tumor progression and metastasis. TSPAN8 may play an important role in mCRC cell invasion. TSPAN8 was overexpressed in human gastric cancer tissues and gastric cancer cell lines compared with the normal. TSPAN8 overexpression promoted cell proliferation and invasion; while TSPAN8 suppression inhibited cell proliferation and invasion. TSPAN8 could activate the ERK MAPK pathway in gastric cancer cells; and MEK-ERK inhibition reversed the effects of TSPAN8 overexpression on cell proliferation and invasion.