A Reliable Research Partner in Life Science and Medicine

Recombinant Human CNPY2 Protein (His Tag)

Catalog No. PKSH030560

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms HP10390;MSAP;TMEM4;ZSIG9

Species Human

HEK293 Cells **Expression Host** Met 1-Ser178 Sequence Accession Q9Y2B0-1 Calculated Molecular Weight 20.0 kDa Observed molecular weight 20 kDa Tag C-His

Bioactivity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CNPY2 is a novel MIR-interacting protein that enhances neurite outgrowth and increases myosin regulatory light chain. CNPY2 enhances migration of C6 glioma cells through phosphorylation of the myosin regulatory light chain. It is

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

expressed in different tissues, including brain. Overexpression of CNPY2 enhanced the motility of glioma cells measured in matrigel invasion chambers and using a scratch assay. Downregulation of CNPY2 by RNA interference significantly decreased glioma cell migration and phosphorylation of MRLC. Inhibition of the corresponding MRLC kinase by ML-7 did not affect migration of CNPY2-overexpressing cells.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com