## Recombinant Human GALNT2/GalNAc-T2 Protein (His Tag)

### Catalog No. PKSH030561

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	GalNAc-T2	
Species	Human	
Expression Host	HEK293 Cells	
Sequence	Lys52-Gln571	
Accession	NP_004472.1	
Calculated Molecular Weight	61.6 kDa	
Observed molecular weight	57-65&46 kDa	
Tag	N-His	
Bioactivity	Not validated for activity	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per $\mu$ g of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from sterile 25 mM Tris, 150 mM NaCl, 0.05% Brij-35, pH 7.5 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	

Data

KDa	MK	R
116	-	
66.2		-
45.0	-	-
35.0	-	
25.0	-	
18.4	-	
14.4	-	

> 95 % as determined by reducing SDS-PAGE.

## Background

GALNT2, also known as GalNAc-T2, is a member of the GalNAc-transferases family. Members of this family transfer an N-acetyl galactosamine to the hydroxyl group of a serine or threonine residue in the first step of O-linked

#### **For Research Use Only**

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

# **Elabscience**®

oligosaccharide biosynthesis. GALNT2 may play a role in lipid metabolism. It catalyzes the initial reaction in O-linked oligosaccharide biosynthesis, the transfer of an N-acetyl-D-galactosamine residue to a serine or threonine residue on the protein receptor. GALNT2 has a broad spectrum of substrates for peptides such as EA2, Muc5AC, Muc1a, Muc1b.