

Recombinant Human BTN3A3 Protein (Fc Tag)

Catalog No. PKSH030635

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Butyrophilin subfamily 3 member A3;BTN3A3;BTF3;BTN3.3
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Trp248
Accession	O00478-1
Calculated Molecular Weight	50.0 kDa
Observed molecular weight	53 kDa
Tag	C-mFc
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Tumor necrosis factor receptor superfamily; member 19 (TNFRSF19); also known as TAJ-alpha or TROY; is a member of the TNF-receptor superfamily. TNFRSF19/TROY expression is detected in the pulmonary epithelium and the ductal

For Research Use Only

epithelium of the prostate and parotid glands. TNFRSF19/TROY expression is detected in some adenocarcinoma cell lines that arise from this tissue. It has been shown to interact with TRAF family members; and to activate JNK signaling pathway when overexpressed in cells. TNFRSF19/TROY is capable of inducing apoptosis by a caspase-independent mechanism; and it is thought to play an essential role in embryonic development. TNFRSF19/TROY was negatively regulated by adipogenic transcription factor CCAAT/enhancer-binding proteins (C/EBP). TNFRSF19 signals activation of the Jnk pathway and induces cell death. Overexpression of TNFRSF19 also signals NFB activation; comparable and similar to that by p75NGFR. TNFRSF19/TROY is capable of activating key signaling pathways of the TNF receptor family; and its predominant expression patterns suggest that it plays a role in the growth and regulation of epithelial tissues.