

Recombinant Human ERP27 Protein (Fc Tag)

Catalog No. PKSH030673

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Endoplasmic Reticulum Resident Protein 27;ER Protein 27;ERp27;ERP27;C12orf46
Species	Human
Expression Host	HEK293 Cells
Sequence	Glu26-Pro269
Accession	Q96DN0
Calculated Molecular Weight	53.7 kDa
Tag	C-mFc
Bioactivity	Not validated for activity

Properties

Purity	> 84 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 84 % as determined by reducing SDS-PAGE.

Background

ERP27 contains 1 thioredoxin domain and is a noncatalytic member of the protein disulfide isomerase family. Protein disulfide isomerases (PDIs) constitute a family of structurally related enzymes which catalyze disulfide bonds formation;

For Research Use Only

reduction; or isomerization of newly synthesized proteins in the lumen of the endoplasmic reticulum (ER). They act also as chaperones; and are; therefore; part of a quality-control system for the correct folding of the proteins in the same subcellular compartment. PDI has been found to have moderate effects (25-fold) on the rate of oxidative folding of proteins in vitro. Recombinant Human Protein Disulfide Isomerase is involved in disulphide-bond formation and isomerization; as well as the reduction of disulphide bonds in proteins. Recombinant PDI has been found to have moderate effects (25-fold) on the rate of oxidative folding of proteins in vitro. ERP27 is a widely expressed protein which localizes to the ER and may act as a protease; protein disulfide isomerase; thiol-disulfide oxidase or phospholipase. ERP27 doesn't contain a CXXC active site motif indicating that it is a catalytically redox-inactive member of the protein disulfide isomerase family.