Recombinant Human WWP2 Protein (His & GST Tag)

Catalog Number:PKSH030722



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms AIP2;WWp2-like

Species Human

Expression Host Baculovirus-Insect Cells

Sequence Met 1-Glu870

Accession O00308

Calculated Molecular Weight 126.7 kDa

Observed molecular weight 120 kDa

Tag N-His-GST

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 8.0, 10% glycerol


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

WWP2 contains 1 C2 domain, 1 HECT (E6AP-type E3 ubiquitin-protein ligase) domain and 4 WW domains. It is an E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. WWP2 can be detected in heart, throughout the brain, placenta, lung, liver, muscle, kidney and pancreas. It is also expressed in spleen and peripheral blood leukocytes. WWP2 polyubiquitinates POU5F1 by 'Lys-63'-linked conjugation and promotes it to proteasomal degradation; in embryonic stem cells (ESCs) the ubiquitination is proposed to regulate POU5F1 protein level. WWP2 ubiquitinates EGR2 and promotes it to proteasomal degradation; in T-cells the ubiquitination inhibits activation-induced cell death. It also ubiquitinates SLC11A2; the ubiquitination is enhanced by presence of NDFIP1 and NDFIP2. WWP2 ubiquitinates RPB1 and promotes

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

Recombinant Human WWP2 Protein (His & GST Tag)

Catalog Number:PKSH030722



it to proteasomal degradation.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com