

Recombinant Human TFAP2C/AP2-GAMMA Protein (His Tag)

Catalog No. PKSH030726

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	AP2-GAMMA;ERF1;hAP-2g;TFAP2G
Species	Human
Expression Host	E.coli
Sequence	Leu128-Val223
Accession	Q92754
Calculated Molecular Weight	12.3 kDa
Observed molecular weight	12-14 kDa
Tag	N-His
Bioactivity	Not validated for activity

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

TFAP2C, also known as AP2-GAMMA, is a member of the activating protein 2 family of transcription factors. AP-2 factors bind to the consensus sequence 5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of important

For Research Use Only

biological functions including proper eye, face, body wall, limb and neural tube development. They also suppress a number of genes including MCAM/MUC18, C/EBP alpha and MYC. TFAP2C may be prognostic indicators for patients with breast tumors. TFAP2C gene has been tested for association to diseases (Breast Neoplasms; Carcinoma) and proposed to participate in processes (cell-cell signaling, male gonad development, regulation of transcription from RNA polymerase II promoter). Proteins are expected to have molecular functions (DNA binding, protein binding, protein dimerization activity, transcription factor activity) and to localize in various compartments (membrane, nucleus).