

Recombinant Human TSPAN1 Protein (aa 110-211, Fc Tag)

Catalog No. PKSH030743

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	NET1;TM4C;TM4SF;TSPAN1
Species	Human
Expression Host	HEK293 Cells
Sequence	Tyr110-Asn211
Accession	O60635
Calculated Molecular Weight	39.1 kDa
Observed molecular weight	43-53 kDa
Tag	N-rFc
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

TSPAN1 belongs to the transmembrane 4 superfamily, also known as the tetraspanin family. Tetraspanins have four hydrophobic domains, intracellular N- and C-termini and two extracellular domains. Tetraspanins act as scaffolding

For Research Use Only

proteins, anchoring multiple proteins to one area of the cell membrane. They also mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. TSPAN1 interacts with human thiamine transporter-1 (hTHTR-1). HTHTR-1 contributes to intestinal thiamine uptake, and its function is regulated at both the transcriptional and posttranscriptional levels. TSPAN1 and hTHTR-1 colocalize in human intestinal epithelial HuTu-80 cells. Coexpression of TSPAN1 in these cells led to a significant decrease in the rate of degradation of hTHTR-1 compared with cells expressing the hTHTR-1 alone; in fact the half-life of the TSPAN1 protein was twice longer in the former cell type compared with the latter cell type.