

Recombinant Human MAD2L1/MAD2 Protein (His Tag)

Catalog Number:PKSH030781



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	HSMAD2;MAD2
Species	Human
Expression Host	E.coli
Sequence	Met 1-Asp 205
Accession	Q13257
Calculated Molecular Weight	25.6 kDa
Observed molecular weight	28 kDa
Tag	N-His

Properties

Purity	> 96 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, 20% glycerol, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 96 % as determined by reducing SDS-PAGE.

Background

Mitotic spindle assembly checkpoint protein MAD2A, also known as HsMAD2, Mitotic arrest deficient 2-like protein 1, MAD2-like protein 1, MAD2L1 and MAD2, is a nucleus and cytoplasm protein which belongs to theMAD2 family. MAD2L1 is a component of the spindle-assembly checkpoint that prevents the onset of anaphase until all chromosomes are properly aligned at the metaphase plate. MAD2L1 is required for the execution of the mitotic checkpoint which monitors the process of kinetochore-spindle attachment and inhibits the activity of the anaphase promoting complex by sequestering CDC20 until all chromosomes are aligned at the metaphase plate. MAD2L1 has two highly different native conformations, an inactive open conformation that cannot bind CDC20 and that predominates in cytosolic monomers, and an active closed conformation. MAD2L1 in the closed conformation preferentially dimerizes with another molecule in the

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human MAD2L1/MAD2 Protein (His Tag)

Catalog Number:PKSH030781



open conformation, but can also form a dimer with a molecule in the closed conformation. Formation of a heterotetrameric core complex containing two molecules of MAD1L1 and of MAD2L1 in the closed conformation promotes binding of another molecule of MAD2L1 in the open conformation and the conversion of the open to the closed form, and thereby promotes interaction with CDC20.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com