

Recombinant Human IFI30 Protein (His Tag)

Catalog No. PKSH030801

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	GILT;IFI-30;IFI30;IP-30;IP30
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Lys243
Accession	P13284
Calculated Molecular Weight	24.7 kDa
Observed molecular weight	31-34 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 96 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 96 % as determined by reducing SDS-PAGE.

Background

IFI30 belongs to the GILT family. This family includes the two characterised human gamma-interferon-inducible lysosomal thiol reductase (GILT) sequences: P13284 and Q9UL08. It also contains several other eukaryotic putative

For Research Use Only

proteins with similarity to GILT. The aligned region contains three conserved cysteine residues. In addition, the two GILT sequences possess a C-X(2)-C motif that is shared by some of the other sequences in the family. This motif is thought to be associated with disulphide bond reduction. IFI30 is a lysosomal thiol reductase that can reduce protein disulfide bonds. It facilitates the generation of MHC class II-restricted epitopes from disulfide bond-containing antigen by the endocytic reduction of disulfide bonds. It also facilitates MHC class I-restricted recognition of exogenous antigens containing disulfide bonds by CD8+ T-cells or crosspresentation. IFI30 may facilitate the complete unfolding of proteins destined for lysosomal degradation and plays an important role in antigen processing.