

## Recombinant Human CD93/C1QR1 Protein (aa 1-580, His Tag)

Catalog No. PKSH030806

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	Complement Component C1q Receptor;C1q/MBL/SPA Receptor;C1qR;C1qR(p);C1qRp;CDw93;Complement Component 1 q Subcomponent Receptor 1;Matrix-Remodeling-Associated Protein 4CD93;C1QR1;MXRA4
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Met 1-Lys 580
<b>Accession</b>	Q9NPY3
<b>Calculated Molecular Weight</b>	59.6 kDa
<b>Observed molecular weight</b>	100 kDa
<b>Tag</b>	C-His
<b>Bioactivity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 90 % as determined by reducing SDS-PAGE.

### For Research Use Only

## Background

CD93 or C1q receptor 1 (C1qR) is an about 120 kDa O-sialoglycoprotein that within the hematopoietic system is selectively expressed on cells of the myeloid lineage. CD93/C1qR is a highly glycosylated transmembrane protein expressed on monocytes; neutrophils; endothelial cells; and stem cells. CD93 was originally identified as a myeloid cell-surface marker and subsequently associated with an ability to modulate phagocytosis of suboptimally opsonized immunoglobulin G and complement particles in vitro. CD93/C1qR; a receptor expressed during early B-cell development; is reinduced during plasma-cell differentiation. High CD93/CD138 expression was restricted to antibody-secreting cells both in T-dependent and T-independent responses as naive; memory; and germinal-center B cells remained CD93-negative. CD93 was expressed on (pre)plasmablasts/plasma cells; including long-lived plasma cells that showed decreased cell cycle activity; high levels of isotype-switched Ig secretion; and modification of the transcriptional network. CD93 is important for the maintenance of plasma cells in bone marrow niches.

## For Research Use Only