Recombinant Human GLT25D2 Protein (His Tag)

Catalog No. PKSH030820

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	C1orf17;GLT25D2;RP11-498P10.2	
Species	Human	
Expression Host	Baculovirus-Insect Cells	
Sequence	Met 1-Ser 622	
Accession	Q8IYK4	
Calculated Molecular Weight	73.8 kDa	
Observed molecular weight	68 kDa	
Tag	C-His	
Bioactivity	Not validated for activity	
Properties		
Purity	> 85 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from sterile 20mM Tris, 500mM NaCl, 10% glycerol, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	
Data		

KDa	MK	R
116 66.2	-	_
45.0	-	
35.0	-	
25.0	-	
18.4 14.4	-	

> 85 % as determined by reducing SDS-PAGE.

Background

Glycosyl transferase 25 domain 2 (GLT25D2) is a glucosyltransferase enzyme expressed only at low levels in the nervous system. Glycosyltransferases are enzymes that act as a catalyst for the transfer of a monosaccharide unit from an activated

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

nucleotide sugar (also known as the "glycosyl donor") to a glycosyl acceptor molecule, usually an alcohol. Glycosyl transferases transfer glycosyl groups onto their substrate. Localization partially defines their function. Glt25D2 enzyme showed a strong galactosyltransferase activity toward various types of collagen and toward the serum mannose-binding lectin MBL which contains a collagen domain.

For Research Use Only