

Recombinant Human DUSP14/MKP-6 Protein (His & MBP Tag)

Catalog No. PKSH030837

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	MKP-L;MKP6
Species	Human
Expression Host	E.coli
Sequence	Met 1 –His 191
Accession	O95147
Calculated Molecular Weight	65.0 kDa
Observed molecular weight	60 kDa
Tag	N-His-MBP
Bioactivity	Not validated for activity

Properties

Purity	> 88 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.5 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 88 % as determined by reducing SDS-PAGE.

Background

Dual specific phosphatase 14 / MAP-kinase phosphatase-6 (DUSP14 / MKP6) is a member of Dual-specificity phosphatases that is a subclass of protein tyrosine phosphatases (PTP) families that can dephosphorylate both

For Research Use Only

phosphotyrosine and phosphoserine / phosphothreonine residues in substrates. Unlike many other DUSPs, DUSP14 only contains a catalytic domain within the C-terminal region. In signal transduction, DUSP14 has been considered as negative regulator of the mitogen-activated protein kinase (MAPK) / extracellular signal-regulated kinase 1 / 2 (ERK 1 / 2) pathway. DUSP14 phosphatase activity has been confirmed to be inhibited by PTP inhibitor IV. PTP inhibitor binds to the catalytic site of DUSP14. PTP inhibitor IV effectively and specifically inhibited DUSP14-mediated dephosphorylation of JNK, a member of the mitogen-activated protein kinase (MAPK) family through dephosphorylation of both the Ser / Thr and Tyr residues of MAPKs.