

Recombinant Human COL9A1 Protein (Fc Tag)

Catalog No. PKSH030873

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	DJ149L1.1.2;EDM6;MED;STL4;Collagen alpha-1(IX) chain;DJ149L1.1.2
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Pro328
Accession	P20849-3
Calculated Molecular Weight	59.2 kDa
Observed molecular weight	62-72 kDa
Tag	C-mFc
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Ubiquinone biosynthesis protein COQ7 homolog; also known as Coenzyme Q biosynthesis protein 7 homolog; Timing protein clk-1 homolog and COQ7; is a mitochondrion inner membrane and peripheral membrane protein which belongs to

For Research Use Only

theCOQ7 family. It is expressed dominantly in heart and skeletal muscle. COQ7 is synthesized as a preprotein that is imported into the mitochondrial matrix; where the sequence is cleaved off and the mature protein becomes loosely associated with the inner membrane. This enzyme is responsible for the hydroxylation of 5-demethoxyubiquinone to 5-hydroxyubiquinone. Human COQ7 protein is mostly helical; and contains an alpha-helical membrane insertion. It has a potential N-glycosylation site; a phosphorylation site for protein kinase C and another for casein kinase II; and three N-myristoylation sites. COQ7 is involved in lifespan determination in ubiquinone-independent manner. It is also involved in ubiquinone biosynthesis. COQ7 is potential central metabolic regulator.