

Recombinant Human Rac2 protein (His tag)

Catalog No. PKSH030903

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	EN-7, Gx, HSPC022, p21-Rac2
Species	Human
Expression Host	E.coli
Sequence	Met 1-Cys 189
Accession	P15153
Calculated Molecular Weight	22.6 kDa
Observed molecular weight	25 kDa
Tag	N-His & C-His
Bioactivity	Testing in progress

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Ras-related C3 botulinum toxin substrate 2 (Rac2) is a small G-protein belonging to the Ras subfamily of the GTPase family. Rac2 acts as an "on / off" switch for signal transduction cascades and motilities. When GDP is attached to the

For Research Use Only

small G-protein, the enzyme is inactivated. Release of the GDP and replace of the GTP cativate the GTPasee. Rac2 remains active until the GTP is hydrolyzed to GDP. Rac2 is a hematopoietic-specific Rho family GTPase implicated as an important constituent of the NADPH oxidase complex and shares 92% amino acid identity with the ubiquitously expressed Rac1. The small G-protein Rac2 regulates the rearrangements of actin and membrane necessary for Fcy receptor-mediated phagocytosis by macrophages. Activated Rac2 binds to the p21-binding domain of PAK1 and this binding provided a basis for microscopic methods to localize activation of these G proteins inside cells.