Recombinant Human XPNPEP2 Protein (His Tag)

Catalog Number: PKSH030976



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Xaa-Pro Aminopeptidase 2;Aminoacylproline Aminopeptidase;Membrane-Bound Aminopeptidase P;Membrane-Bound APP;Membrane-Bound AmP;mAmP;X-Pro Aminopeptidase 2;XPNPEP2
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Ala 650
Accession	O43895
Calculated Molecular Weight	72.0 kDa
Tag	C-His
Bioactivity	Measured by its ability to cleave the fluorogenic peptide substrate, H- Lys(2-Aminobenzoyl)Pro-Pro-pNitroanilide(K(Abz)PP-pNA). The specific activity is > 300 pmoles/min/µg.
Properties	
Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

KDa 116 66.2	MK R
45.0 35.0	-
25.0	-
18.4 14.4	=

> 97 % as determined by reducing SDS-PAGE.

Background

Aminopeptidase P (APP) is a hydrolase specific for N-terminal imido bonds; which are common to several collagen degradation products; neuropeptides; vasoactive peptides; and cytokines. A membrane-bound and soluble form of this enzyme (XPNPEP2) have been identified as products of two separate genes. XPNPEP2; the X-linked gene that encodes membranous aminopeptidase P (APP); has been reported to associate with APP activity. The membrane aminopeptidase

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Catalog Number: PKSH030976



P (XPNPEP2) is largely limited in distribution to endothelia and brush border epithelia. APP and XPNPEP2 contain homologous blocks of sequence common to members of the "pita bread-fold" protein family; of which Escherichia coli methionine aminopeptidase is the prototype. The C-2399A variant in XPNPEP2 is associated with reduced APP activity and a higher incidence of AE-ACEi. XPNPEP2 mRNA was detected in fibroblasts that carry the translocation; suggesting that this gene at least partially escapes X inactivation. XPNPEP2 is a candidate gene for premature ovarian failure (POF).

For Research Use Only

A Reliable Research Partner in Life Science and Medicine
Toll-free: 1-888-852-8623 Tel: 1-832-243-6086
Web: www.elabscience.com Email: techsupport@elabscience.com