

## Recombinant Human NRXN3 Protein (His Tag)

**Catalog No.** PKSH030984

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	C14orf60
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Met 1-Thr 357
<b>Accession</b>	NP_620426.2
<b>Calculated Molecular Weight</b>	36.0 kDa
<b>Observed molecular weight</b>	50-60 kDa
<b>Tag</b>	C-His
<b>Bioactivity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 94 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 94 % as determined by reducing SDS-PAGE.

### Background

Neurexin-3-beta; also known as Neurexin III-beta and NRXN3; is a single-pass type I membrane protein which belongs to the neurexin family. It contains onelaminin G-like domain. It is a neuronal cell surface protein that may be involved in cell

### For Research Use Only

recognition and cell adhesion. Neurexins are a family of proteins that function in the vertebrate nervous system as cell adhesion molecules and receptors. They are encoded by several unlinked genes of which two; NRXN1 and NRXN3; are among the largest known human genes. Three of the genes ( NRXN1; NRXN2; NRXN3 ) utilize two alternate promoters and include numerous alternatively spliced exons to generate thousands of distinct mRNA transcripts and protein isoforms. The majority of transcripts are produced from the upstream promoter and encode alpha-neurexin isoforms; a much smaller number of transcripts are produced from the downstream promoter and encode beta-neurexin isoforms. The alpha-neurexins contain EGF-like sequences and laminin G domains; and have been shown to interact with neurexophilins. The beta-neurexins lack EGF-like sequences and contain fewer laminin G domains than alpha-neurexins. NRXN3 have been linked to genetic predisposition towards a number of conditions such as alcohol or drug addiction; or obesity.