A Reliable Research Partner in Life Science and Medicine

Recombinant Human CD1B/CD1A Protein (Fc Tag)

Catalog No. PKSH030990

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms CD1;CD1A;CD1B;R1

Species Human

Expression Host HEK293 Cells
Sequence Met 1-Ser 303
Accession NP_001755
Calculated Molecular Weight 58.6 kDa
Observed molecular weight 59-69 kDa
Tag C-hFc

Bioactivity Not validated for activity

Properties

Purity > 88 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 88 % as determined by reducing SDS-PAGE.

Background

CD1B contains 1 Ig-like (immunoglobulin-like) domain and belongs to the CD1 family. CD1 family members are transmembrane glycoproteins, which are structurally related to the major histocompatibility complex (MHC) proteins and

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

form heterodimers with beta-2-microglobulin. During protein synthesis and maturation, they bind endogenous lipids that are replaced by lipid or glycolipid antigens when the proteins are internalized and pass through endosomes, before trafficking back to the cell surface. CD1B localizes to late endosomes and lysosomes via a tyrosine-based motif in the cytoplasmic tail, and requires vesicular acidification to bind lipid antigens.. It is expressed on cortical thymocytes, epidermal Langerhans cells, dendritic cells, on certain T-cell leukemias, and in various other tissues. CD1B is an antigenpresenting protein that binds self and non-self lipid and glycolipid antigens and presents them to T-cell receptors on natural killer T-cells.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com