

Recombinant Human HDAC4 Protein (aa 612-1084)

Catalog Number:PKSH030998



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	AHO3;BDMR;HA6116;HD4;HDAC-4;HDAC-A;HDACA
Species	Human
Expression Host	Baculovirus-Insect Cells
Sequence	Met612-Leu1084
Accession	NP_006028.2
Calculated Molecular Weight	50.9 kDa
Observed molecular weight	51 kDa
Tag	None

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20mM Tris,500mM NaCl, pH 7.4, 10% glycerol Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

HDAC4 (histone deacetylase 4), belongs to class II of the histone deacetylase/acuc/apha family. Histone Deacetylases (HDACs) are a group of enzymes closely related to sirtuins. They catalyze the removal of acetyl groups from lysine residues in histones and non-histone proteins, resulting in transcriptional repression. In general, they do not act autonomously but as components of large multiprotein complexes, such as pRb-E2F and mSin3A, that mediate important transcription regulatory pathways. There are three classes of HDACs; classes 1, 2 and 4, which are closely related Zn²⁺-dependent enzymes. HDACs are ubiquitously expressed and they can exist in the nucleus or cytosol. Their subcellular localization is effected by protein-protein interactions and by the class to which they belong. HDACs have a role in cell growth arrest, differentiation and death and this has led to substantial interest in HDAC inhibitors as possible

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human HDAC4 Protein (aa 612-1084)

Catalog Number:PKSH030998



antineoplastic agents. HDAC4 possesses histone deacetylase activity and represses transcription when tethered to a promoter. It does not bind DNA directly, but through transcription factors MEF2C and MEF2D. HDAC4 seems to interact in a multiprotein complex with RbAp48 and HDAC3.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com