

# Recombinant Human CD200R1L/CD200R2/CD200RLa Protein (His Tag)



Catalog Number:PKSH031061

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	CD200R2;CD200RLa
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Met 1-Leu239
<b>Accession</b>	AAT00538.1
<b>Calculated Molecular Weight</b>	25.2 kDa
<b>Observed molecular weight</b>	45-60 kDa
<b>Tag</b>	C-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Cell surface glycoprotein CD200 receptor 2, also known as Cell surface glycoprotein CD200 receptor 1-like, Cell surface glycoprotein OX2 receptor 2, CD200 receptor-like 2, CD200R1a, CD200R1L and CD200R2, is a single-pass type I membrane protein which belongs to the CD200R family. CD200R1L / CD200R2. It contains one Ig-like C2-type (immunoglobulin-like) domain and one Ig-like V-type (immunoglobulin-like) domain. CD200 is a transmembrane protein delivering immunoregulatory signals after engagement of CD200R. A family of CD200Rs exist ( CD200R1, CD200R2, CD200R3, CD200R4 ) with different tissue expression and functional activity. In the presence of anti-CD200R2 / CD200R3 monoclonal antibodies (mAbs), bone-marrow cells cultured in the presence of (interleukin [IL]-4+granulocyte-macrophage colony-stimulating factor) differentiate into dendritic cells (DCs), which induce CD4+CD25+ Treg.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human CD200R1L/CD200R2/CD200RLa Protein (His Tag)



Catalog Number:PKSH031061

---

Interaction between the relatively ubiquitously expressed molecule CD200 and one of its receptors, CD200R1, resulted in direct suppression of alloreactivity, engagement of alternate receptors led instead to altered differentiation of dendritic cells (DCs) from marrow precursors, which could in turn foster development of Foxp3(+) regulatory T cells. Unlike anti-CD200R1, anti-CD200R2 both promotes development of DCs with capacity to induce Treg and directly augments thymocyte production of Treg.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)