

Recombinant Human PTPN12 Protein (aa 1-355, His & GST Tag)

Catalog No. PKSH031089

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	PTP-PEST;PTPG1
Species	Human
Expression Host	Baculovirus-Insect Cells
Sequence	Met 1-Gln355
Accession	AAA36529.1
Calculated Molecular Weight	69.4 kDa
Observed molecular weight	64 kDa
Tag	N-His-GST
Bioactivity	Measured by its ability to dephosphorylate a phosphotyrosine residue in an EGF receptor 988-998 phosphopeptide substrate, R&D Systems, Catalog # ES006. The specific activity is > 15 μ moles/min/mg.

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20mM Tris, 500mM NaCl, 10% glycerol, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

PTPN12 is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. PTPN12 contains a C-terminal PEST motif, which serves as a protein–protein interaction domain, and may be related to protein intracellular half-life. PTPN12 was found to bind and dephosphorylate the product of oncogene c-ABL, thus may play a role in oncogenesis. PTPN12 was shown to interact with, and dephosphorylate, various of cytoskeleton and cell adhesion molecules, such as p130 (Cas), CAKbeta/PTK2B, PSTPIP1, and paxillin, which suggested its regulatory roles in controlling cell shape and mobility.