Recombinant Human HDAC3 Protein (His & GST Tag)

Catalog No. PKSH031105

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	HD3;RPD3;RPD3-2
Species	Human
Expression Host	Baculovirus-Insect Cells
Sequence	Met 1-Ile 428
Accession	O15379-1
Calculated Molecular Weight	76.7 kDa
Observed molecular weight	66 kDa
Tag	N-His-GST
Bioactivity	Not validated for activity
Properties	
Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20mM Tris, 500mM NaCl, 4mM GSH, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Histone Deacetylases (HDACs) are a group of enzymes closely related to sirtuins. They catalyze the removal of acetyl groups from lysine residues in histones and non-histone proteins, resulting in transcriptional repression. In general, they do

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

not act autonomously but as components of large multiprotein complexes, such as pRb-E2F and mSin3A, that mediate important transcription regulatory pathways. There are three classes of HDACs, classes 1, 2 and 4, which are closely related Zn2+-dependent enzymes. HDACs are ubiquitously expressed and they can exist in the nucleus or cytosol. Their subcellular localization is effected by protein-protein interactions and by the class to which they belong. HDACs have a role in cell growth arrest, differentiation and death and this has led to substantial interest in HDAC inhibitors as possible antineoplastic agents. Histone Deacetylases (HDACs) is Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Probably participates in the regulation of transcription through its binding to the zinc-finger transcription factor YY1, increases YY1 repression activity. Required to repress transcription of the POU1F1 transcription factor. Acts as a molecular chaperone for shuttling phosphorylated NR2C1 to PML bodies for sumoylation.

For Research Use Only